

Upper Owyhee Watershed Assessment

Appendix J. Riparian plant associations

© Owyhee Watershed Council and Scientific Ecological Services

Contents

Inter-mountain basins semi-desert grassland.
Great Basin Foothill and Lower Montane Riparian Woodland and Shrubland
Columbia Basin Foothill Riparian Woodland and Shrubland
Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland
Rocky Mountain Subalpine-Montane Riparian Shrubland
Rocky Mountain Subalpine-Montane Riparian Woodland
Columbia Plateau Silver Sagebrush Seasonally Flooded Shrub-Steppe
North American Arid West Emergent Marsh
Rocky Mountain Subalpine-Montane Mesic Meadow
Rocky Mountain Subalpine Mesic Meadow
Rocky Mountain Alpine-Montane Wet Meadow

Riparian plant associations

Inter-mountain basins semi-desert grassland.

This widespread ecological system occurs throughout the Intermountain western U.S. on dry plains and mesas, at approximately 1450 to 2320 m (4750-7610 feet) in elevation. These grasslands occur in lowland and upland areas and may occupy swales, playas, mesa tops, plateau parks, alluvial flats, and plains, but sites are typically xeric. Substrates are often well-drained sandy- or loamy-textured soils derived from sedimentary parent materials, but are quite variable and may include fine-textured soils derived from igneous and metamorphic rocks. When they occur near foothills grasslands they will be at lower elevations. The dominant perennial bunch grasses and shrubs within this system are all very drought-resistant plants. These grasslands are typically dominated or codominated by *Achnatherum hymenoides*, *Aristida* spp., *Bouteloua gracilis*, *Hesperostipa comata*, *Muhlenbergia torreyana*, or *Pleuraphis jamesii*, and may include scattered shrubs and dwarf-shrubs of species of *Artemisia*, *Atriplex*, *Coleogyne*, *Ephedra*, *Gutierrezia*, or *Krascheninnikovia lanata*.

Great Basin Foothill and Lower Montane Riparian Woodland and Shrubland

This system occurs in mountain ranges of the Great Basin and along the eastern slope of the Sierra Nevada within a broad elevation range from about 1220 m (4000

feet) to over 2135 m (7000 feet). This system often occurs as a mosaic of multiple communities that are tree-dominated with a diverse shrub component. The variety of plant associations connected to this system reflects elevation, stream gradient, floodplain width, and flooding events. Dominant trees may include *Abies concolor*, *Alnus incana*, *Betula occidentalis*, *Populus angustifolia*, *Populus balsamifera* ssp. *trichocarpa*, *Populus fremontii*, *Salix laevigata*, *Salix gooddingii*, and *Pseudotsuga menziesii*. Dominant shrubs include *Artemisia cana*, *Cornus sericea*, *Salix exigua*, *Salix lasiolepis*, *Salix lemmonii*, or *Salix lutea*. Herbaceous layers are often dominated by species of *Carex* and *Juncus*, and perennial grasses and mesic forbs such *Deschampsia caespitosa*, *Elymus trachycaulus*, *Glyceria striata*, *Iris missouriensis*, *Maianthemum stellatum*, or *Thalictrum fendleri*. Introduced forage species such as *Agrostis stolonifera*, *Poa pratensis*, *Phleum pratense*, and the weedy annual *Bromus tectorum* are often present in disturbed stands. These are disturbance-driven systems that require flooding, scour and deposition for germination and maintenance. Livestock grazing is a major influence in altering structure, composition, and function of the community.

Columbia Basin Foothill Riparian Woodland and Shrubland

This is a low-elevation riparian system found on the periphery of the mountains surrounding the Columbia River Basin, along major tributaries and the main stem of the Columbia at relatively low elevations. This is the riparian system associated with all streams at and below lower treeline, including permanent, intermittent and ephemeral streams with woody riparian vegetation. These forests and woodlands require flooding and some gravels for reestablishment. They are found in low-elevation canyons and draws, on floodplains, or in steep-sided canyons, or narrow V-shaped valleys with rocky substrates. Sites are subject to temporary flooding during spring runoff. Underlying gravels may keep the water table just below the ground surface and are favored substrates for cottonwood. Large bottomlands may have large occurrences, but most have been cut over or cleared for agriculture. Rafted ice and logs in freshets may cause considerable damage to tree boles. Beavers crop younger cottonwood and willows and frequently dam side channels occurring in these stands. In steep-sided canyons, streams typically have perennial flow on mid to high gradients. Important and diagnostic trees include *Populus balsamifera* ssp. *trichocarpa*, *Alnus rhombifolia*, *Populus tremuloides*, *Celtis laevigata* var. *reticulata*, *Betula occidentalis*, or *Pinus ponderosa*. Important shrubs include *Crataegus douglasii*, *Philadelphus lewisii*, *Cornus sericea*, *Salix lucida* ssp. *lasiandra*, *Salix eriocephala*, *Rosa nutkana*, *Rosa woodsii*, *Amelanchier alnifolia*, *Prunus virginiana*, and *Symphoricarpos albus*. Grazing is a major influence in altering structure, composition, and function of the community.

Rocky Mountain Lower Montane-Foothill Riparian Woodland and Shrubland

This ecological system is found throughout the Rocky Mountain and Colorado Plateau regions within a broad elevational range from approximately 900 to 2800 m. This system often occurs as a mosaic of multiple communities that are tree-dominated with a diverse shrub component. It is dependent on a natural hydrologic regime, especially annual to episodic flooding. Occurrences are found within the flood zone of

rivers, on islands, sand or cobble bars, and immediate streambanks. It can form large, wide occurrences on mid-channel islands in larger rivers or narrow bands on small, rocky canyon tributaries and well-drained benches. It is also typically found in backwater channels and other perennially wet but less scoured sites, such as floodplains swales and irrigation ditches. In some locations, occurrences extend into moderately high intermountain basins where the adjacent vegetation is sage steppe. Dominant trees may include *Acer negundo*, *Populus angustifolia*, *Populus deltoides*, *Populus fremontii*, *Pseudotsuga menziesii*, *Picea pungens*, *Salix amygdaloides*, or *Juniperus scopulorum*. Dominant shrubs include *Acer glabrum*, *Alnus incana*, *Betula occidentalis*, *Cornus sericea*, *Crataegus rivularis*, *Forestiera pubescens*, *Prunus virginiana*, *Rhus trilobata*, *Salix monticola*, *Salix drummondiana*, *Salix exigua*, *Salix irrorata*, *Salix lucida*, *Shepherdia argentea*, or *Symphoricarpos* spp. Exotic trees of *Elaeagnus angustifolia* and *Tamarix* spp. are common in some stands. Generally, the upland vegetation surrounding this riparian system is different and ranges from grasslands to forests. In the Wyoming Basins, the high-elevation *Populus angustifolia*-dominated rivers are included here, including along the North Platte, Sweetwater, and Laramie rivers. In these situations, *Populus angustifolia* is extending down into the sage steppe zone of the basins.

Rocky Mountain Subalpine-Montane Riparian Shrubland

This system is found throughout the Rocky Mountain cordillera from New Mexico north into Montana, and also occurs in mountainous areas of the Intermountain region and Colorado Plateau. These are montane to subalpine riparian shrublands occurring as narrow bands of shrubs lining streambanks and alluvial terraces in narrow to wide, low-gradient valley bottoms and floodplains with sinuous stream channels. Generally it is found at higher elevations, but can be found anywhere from 1700-3475 m. Occurrences can also be found around seeps, fens, and isolated springs on hillslopes away from valley bottoms. Many of the plant associations found within this system are associated with beaver activity. This system often occurs as a mosaic of multiple communities that are shrub- and herb-dominated and includes above-treeline, willow-dominated, snowmelt-fed basins that feed into streams. The dominant shrubs reflect the large elevational gradient and include *Alnus incana*, *Betula nana*, *Betula occidentalis*, *Cornus sericea*, *Salix bebbiana*, *Salix boothii*, *Salix brachycarpa*, *Salix drummondiana*, *Salix eriocephala*, *Salix geyeriana*, *Salix monticola*, *Salix planifolia*, and *Salix wolfii*. Generally the upland vegetation surrounding these riparian systems are of either conifer or aspen forests.

Rocky Mountain Subalpine-Montane Riparian Woodland

This riparian woodland system is comprised of seasonally flooded forests and woodlands found at montane to subalpine elevations of the Rocky Mountain cordillera, from southern New Mexico north into Montana, and west into the Intermountain region and the Colorado Plateau. It occurs throughout the interior of British Columbia and the eastern slopes of the Cascade Mountains. This system contains the conifer and aspen woodlands that line montane streams. These are communities tolerant of periodic flooding and high water tables. Snowmelt moisture in this system may create shallow water tables or seeps for a portion of the growing season. Stands typically occur at

elevations between 1500 and 3300 m (4920-10,830 feet), farther north elevation ranges between 900 and 2000 m. This is confined to specific riparian environments occurring on floodplains or terraces of rivers and streams, in V-shaped, narrow valleys and canyons (where there is cold-air drainage). Less frequently, occurrences are found in moderate-wide valley bottoms on large floodplains along broad, meandering rivers, and on pond or lake margins. Dominant tree species vary across the latitudinal range, although it usually includes *Abies lasiocarpa* and/or *Picea engelmannii*; other important species include *Pseudotsuga menziesii*, *Picea pungens*, *Picea engelmannii* X *glauca*, *Populus tremuloides*, and *Juniperus scopulorum*. Other trees possibly present but not usually dominant include *Alnus incana*, *Abies concolor*, *Abies grandis*, *Pinus contorta*, *Populus angustifolia*, *Populus balsamifera* ssp. *trichocarpa*, and *Juniperus osteosperma*.

Columbia Plateau Silver Sagebrush Seasonally Flooded Shrub-Steppe

This ecological system includes sagebrush communities occurring at lowland and montane elevations in the Columbia Plateau-northern Great Basin region, east almost to the Great Plains. These are generally depressional wetlands or non-alkaline playas, occurring as small- or occasionally large-patch communities, in a sagebrush or montane forest matrix. Climate is generally semi-arid, although it can be cool in montane areas. This system occurs in poorly drained depressional wetlands, the largest characterized as playas, the smaller as vernal pools, or along seasonal stream channels in valley bottoms or mountain meadows. *Artemisia cana* ssp. *bolanderi* or *Artemisia cana* ssp. *viscidula* are dominant, with *Artemisia tridentata* ssp. *tridentata*, *Artemisia tridentata* ssp. *wyomingensis*, or *Artemisia tridentata* ssp. *vaseyana* occasionally codominant. Understory graminoids and forbs are characteristic, with *Poa secunda* (= *Poa nevadensis*), *Poa cusickii*, *Muhlenbergia filiformis*, *Muhlenbergia richardsonis*, and *Leymus cinereus* dominant at the drier sites; *Eleocharis palustris*, *Deschampsia caespitosa*, and *Carex* species dominate at wetter or higher-elevation sites.

North American Arid West Emergent Marsh

This widespread ecological system occurs throughout much of the arid and semi-arid regions of western North America, typically surrounded by savanna, shrub steppe, steppe, or desert vegetation. Natural marshes may occur in depressions in the landscape (ponds, kettle ponds), as fringes around lakes, and along slow-flowing streams and rivers (such riparian marshes are also referred to as sloughs). Marshes are frequently or continually inundated, with water depths up to 2 m. Water levels may be stable, or may fluctuate 1 m or more over the course of the growing season. Water chemistry may include some alkaline or semi-alkaline situations, but the alkalinity is highly variable even within the same complex of wetlands. Marshes have distinctive soils that are typically mineral, but can also accumulate organic material. Soils have characteristics that result from long periods of anaerobic conditions in the soils (e.g., gleyed soils, high organic content, redoximorphic features). The vegetation is characterized by herbaceous plants that are adapted to saturated soil conditions. Common emergent and floating vegetation includes species of *Scirpus* and/or *Schoenoplectus*, *Typha*, *Juncus*, *Potamogeton*, *Polygonum*, *Nuphar*, and *Phalaris*. This system may also include areas of relatively deep water with floating-leaved plants

(*Lemna*, *Potamogeton*, and *Brasenia*) and submergent and floating plants (*Myriophyllum*, *Ceratophyllum*, and *Elodea*).

Rocky Mountain Subalpine-Montane Mesic Meadow

This Rocky Mountain ecological system is restricted to sites from lower montane to subalpine where finely textured soils, snow deposition, or windswept dry conditions limit tree establishment. It is found typically above 2000 m in elevation in the southern part of its range and above 600 m in the northern part. These upland communities occur on gentle to moderate-gradient slopes. The soils are typically seasonally moist to saturated in the spring, but if so will dry out later in the growing season. These sites are not as wet as those found in Rocky Mountain Alpine-Montane Wet Meadow (CES306.812). Vegetation is typically forb-rich, with forbs contributing more to overall herbaceous cover than graminoids. Important taxa include *Erigeron* spp., *Asteraceae* spp., *Mertensia* spp., *Penstemon* spp., *Campanula* spp., *Lupinus* spp., *Solidago* spp., *Ligusticum* spp., *Thalictrum occidentale*, *Valeriana sitchensis*, *Rudbeckia occidentalis*, *Balsamorhiza sagittata*, *Wyethia* spp., *Deschampsia caespitosa*, *Koeleria macrantha*, and *Dasiphora fruticosa*. Burrowing mammals can increase the forb diversity.

Rocky Mountain Subalpine Mesic Meadow

This Rocky Mountain ecological system is restricted to sites in the subalpine zone where finely textured soils, snow deposition, or wind-swept dry conditions limit tree establishment. It is found typically above 3000 m in elevation in the southern part of its range and above 1500 m in the northern part. These upland communities occur on gentle to moderate-gradient slopes. The soils are typically seasonally moist to saturated in the spring, but if so will dry out later in the growing season. These sites are not as wet as found in Rocky Mountain Alpine-Montane Wet Meadow (CES306.812). Vegetation is typically forb-rich, with forbs contributing more to overall herbaceous cover than graminoids. Important taxa include *Erigeron* spp., *Asteraceae* spp., *Mertensia* spp., *Penstemon* spp., *Campanula* spp., *Lupinus* spp., *Solidago* spp., *Ligusticum* spp., *Thalictrum occidentale*, *Valeriana sitchensis*, *Balsamorhiza sagittata*, *Wyethia* spp., *Deschampsia caespitosa*, *Koeleria macrantha*, and *Dasiphora fruticosa*. Burrowing mammals can increase the forb diversity.

Rocky Mountain Alpine-Montane Wet Meadow

These are high-elevation communities found throughout the Rocky Mountains and Intermountain regions, dominated by herbaceous species found on wetter sites with very low-velocity surface and subsurface flows. They range in elevation from montane to alpine (1000-3600 m). These types occur as large meadows in montane or subalpine valleys, as narrow strips bordering ponds, lakes, and streams, and along toeslope seeps. They are typically found on flat areas or gentle slopes, but may also occur on sub-irrigated sites with slopes up to 10%. In alpine regions, sites typically are small depressions located below late-melting snow patches or on snowbeds. Soils of this system may be mineral or organic. In either case, soils show typical hydric soil characteristics, including high organic content and/or low chroma and redoximorphic features. This system often occurs as a mosaic of several plant associations, often

dominated by graminoids, including *Calamagrostis stricta*, *Caltha leptosepala*, *Cardamine cordifolia*, *Carex illota*, *Carex microptera*, *Carex nigricans*, *Carex scopulorum*, *Carex utriculata*, *Carex vernacula*, *Deschampsia caespitosa*, *Eleocharis quinqueflora*, *Juncus drummondii*, *Phippsia algida*, *Rorippa alpina*, *Senecio triangularis*, *Trifolium parryi*, and *Trollius laxus*. Often alpine dwarfshrublands, especially those dominated by *Salix*, are immediately adjacent to the wet meadows. Wet meadows are tightly associated with snowmelt and typically not subjected to high disturbance events such as flooding.

Bibliography

- Nature Serve. 2003. Appendix E - Landcover descriptions for the Southwest regional GAP analysis project. Retrieved 7/11/2010. <http://www.waterresources.slco.org/pdf/WaQSPAppE.pdf>
- Nature Serve. 2005. Land cover legend: Current distribution of sagebrush and associated vegetation in the Columbia Basin and southwestern regions. Accessed 1/2/2010. http://sagemap.wr.usgs.gov/FTP/Documents/Shrubmap_Legend_Decriptions.pdf
- Oregon Department of Forestry. 995. AppendixD: Plant associations. In *Eastern Region Long-Range Forest Management Plan*. Accessed 1/7/2011. http://www.oregon.gov/ODF/STATE_FORESTS/docs/management/eor/D_pl_asc.pdf?ga=t
- Oregon Wildlife Explorer. 2008. Wetland and riparian habitats. Accessed 1/2/2011. <http://www.oregonexplorer.info/wildlife/WildlifeHabitatClassification/WetlandandRiparian>
- USGS National Biological Information Infrastructure. Gap analysis program northwest. Accessed 7/11/2010. <http://gap.uidaho.edu/index.php/gap-home/Northwest-GAP/landcover>
- USGS National Biological Information Infrastructure. Welcome to the GAP analysis program (GAP) land cover viewer. Accessed 7/11/2010. <http://lc.gapanalysisprogram.com/landcoverviewer/>